

2015-12

Permanent draft genome of 'Rhodopirellula islandica' strain K833.

Kizina, J

<http://hdl.handle.net/10026.1/9388>

10.1016/j.margen.2015.07.011

Mar Genomics

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **Marine Genomics – Genomics/Technical resources**

2

3 **Permanent draft genome of ‘*Rhodopirellula islandica*’ strain K833**

4

5 Jana Kizina¹, Marina Zure¹, Colin Bernhard Munn³, Michael Richter², Jens Harder¹

6

7 ¹ Max Planck Institute for Marine Microbiology, Department of Microbiology, D-28359 Bremen,
8 Germany, ² Max Planck Institute for Marine Microbiology, Microbial Genomics and Bioinformatics
9 Research Group, D-28359 Bremen, Germany, ³School of Marine Sciences and Engineering,
10 University of Plymouth, Plymouth PL4 8AA, United Kingdom

11

12 Corresponding author: Jens Harder, E-Mail: jharder@mpi-bremen.de, Phone: +49 (0)421/2028750,
13 Fax: +49 (0)421/2028580

14

15 Competing interest: The authors have declared that no competing interests exist.

16

17 **Keywords**

18 *Planctomycetes*, cold adaptation

19

20 **Abstract**

21 **The ‘*Rhodopirellula islandica*’ strain K833 has a lower temperature range for growth than other**

22 genome-sequenced *Rhodopirellula baltica* and '*Rhodopirellula europaea*' strains. The draft
23 genome of K833 was obtained as part of a larger study on the biogeography of *Rhodopirellula*
24 species in European marine waters. The genome consists of 70 contigs with a genome size of
25 7439654 bp. With an average nucleotide identity of XXX % to related genomes of
26 *Rhodopirellula baltica* and '*Rhodopirellula europaea*', it will be a valuable source for the study
27 of temperature adaptation of planctomycetes.

28

29 Introduction

30 *Rhodopirellula* is a genus of marine *Planctomycetes* which are unusual bacteria lacking a
31 peptidoglycan membrane. *Planctomycetes* live frequently attached to surfaces, they are abundant in
32 the particulate fractions of marine ecosystems and considered as important participants in the global
33 carbon and nitrogen cycles. *Rhodopirellula baltica* SH1(T) was the source for the first planctomycete
34 genome (Glöckner et al. 2003). A collection of 70 *Rhodopirellula* strains obtained from different
35 European seas (Winkelmann and Harder, 2009) revealed 13 distinct operational taxonomic units
36 (OTUs). (Winkelmann et al., 2010). Eight strains were sequenced and covered sample sites from the
37 Baltic Sea to the Mediterranean Sea (Klindworth et al., 2014, Richter et al., 2014a,b, Richter-
38 Heitmann et al., 2014, Wegner et al., 2014). '*Rhodopirellula islandica*' strain K833 (= JCM 17612 =
39 DSM 24040) was isolated from a water sample on the coast of Sandgerdi, Island (64.0356 N 22.6986
40 W) (Winkelmann and Harder, 2009). In a comparative growth study with cells grown on M13a plates
41 in 250ml flasks that contained 50 ml of M13 medium in artificial seawater at ~ 35 per mille salinity
42 and were shaken at 50 rpm in the dark at 7, 14, 21, 28 or 35 °C, K833 grew in contrast to the other
43 strains not at 35 °C, whereas *Rhodopirellula baltica* SH1(T) grew and '*Rhodopirellula europaea*' 6C
44 had the highest growth rate at 35 °C. At 14 °C, 6C had a long lag phase before growth started and
45 K833 had of the three strains the fastest growth rate. Thus, the strains are a valuable source for the
46 investigation of temperature adaptation of microorganisms. The average nucleotide identity between
47 these strains is XXXX, indicating a very close relationship between the three species.

48 Data description

49 Genomic DNA of K833 was sequenced by the Illumina MiSeq technology at the Max Planck-
50 Genome-centre Cologne. 5494521 paired-end reads of 250 bp were dynamically trimmed with
51 SolexaQA v.2.2. (Cox et al. 2010) and normalized with khmer 1.0 (Crusoe et al. 2014). 1462500
52 high-quality reads were assembled with Spades 3.1.0linux (Bankevich 2012). Contigs were *de-novo*
53 assembled in Geneious R8 (Biomatters, Auckland, New Zealand) to remove duplications and reads
54 were mapped with BBtools to identify possible contig elongations. The mapping reads were
55 reassembled using the first assembly as trusted assembly in Spades. After six rounds, the assembly
56 was stable and CheckM 0.9 indicated a completeness of 99.93% with a contamination value of 0.0 %
57 (Parks et al. 2014). The genome was annotated in RAST (Aziz et al. 2008).

58 Michael – ergänzungen - literatur

59 The genome encodes XXXX proteins, xxx tRNAs and xxx rRNAs. These values are in the range of
60 previously reported *Rhodopirellula* strains, with over 7 Mb and 6000 predicted open reading frames
61 each, and reflects the complex lifestyle of the planctomycetes. Pairwise analysis by reciprocal best
62 match BLAST revealed 4241 shared genes between the strain and *R. baltica* SH1^T. This high number
63 reflects the close relation between the two species as predicted by 16S rDNA and ANI analysis. The
64 sessile lifestyle of planctomycetes comprises life in oxygen-limited biofilms. The K833 genome
65 codes for the synthesis of menaquinon, the typical quinon of microaerophiles, and a menaquinon
66 (vitamin K)-dependent gamma-carboxylase that is not present in *R. baltica* SH1^T. The genome of
67 K833 codes not only for cbb₃-type cytochrome c and cytochrome d oxidases – well known for their
68 high affinity to molecular oxygen -, but also for a periplasmatic nitrate reduction pathway as
69 alternative electron acceptor which is lacking in *R. baltica* SH1^T. Thus, the bacterium is well adapted
70 to microoxic-anoxic transition zones. A feature of these transition zones is fermentation yielding an
71 acidification. In contrast to the *R. baltica* SH1^T, the K833 genome contains a glutamate decarboxylase
72 (EC 4.1.1.15) conferring acid-resistance. It has also some unique sulfatases and glycosyl hydrolases.
73 Unusual is the presence of *traG* and *traI* genes within a region of hypothetical proteins, indicating a

74 potential for genetic transfer.

75

76 **Nucleotide sequence accession number.** The Whole Genome Shotgun projects was deposited in
77 INSDC (DDBJ/EBI-ENA/GenBank) under the accession number XXXXXXXX.

78

79 **Acknowledgments**

80 This study was funded by the Max Planck Society. Marina Zure was supported through a MARES
81 Grant. MARES is a Joint Doctorate programme selected under Erasmus Mundus coordinated by
82 Ghent University (FPA 2011-0016). See www.mares-eu.org for extra information.

83

84 **References**

85 Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A., Formsma, K., Gerdes, S.,
86 Glass, E.M., Kubal, M., Meyer, F., Olsen, G.J., Olson, R., Osterman, A.L., Overbeek, R.A., McNeil,
87 L.K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G.D., Reich, C., Stevens, R., Vassieva, O.,
88 Vonstein, V., Wilke, A., Zagnitko, O. (2008) The RAST Server: rapid annotations using subsystems
89 technology. BMC Genomics 9,75.

90 Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
91 Nikolenko, S.I., Pham, S., Prjibelski, A.D. Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G.,
92 Alekseyev, M.A., Pevzner P.A. (2012) SPAdes: A new genome assembly algorithm and its
93 applications to single-cell sequencing. J. Comput. Biol. 19, 455–477.

94 Cox, M.P., Peterson, D.A., Biggs, P.J. (2010) SolexaQA: At-a-glance quality assessment of Illumina
95 second-generation sequencing data. BMC Bioinformatics 11:485.

96 Crusoe, M., Edverson, G., Fish, J., Howe, A., McDonald, E., Nahum, J., Nanlohy, K., Ortiz-
 97 Zuazaga, H., Pell, J., Simpson, J., Scott, C., Srinivasan, R.R., Zhang, Q., Brown, C. T. (2014) The
 98 khmer software package: enabling efficient sequence analysis. Figshare
 99 <http://dx.doi.org/10.6084/m9.figshare.979190>
 100
 101 Glöckner, F. O., Kube, M., Bauer, M., Teeling, H., Lombardot, T., Ludwig, W., Gade, D., Beck, A.,
 102 Borzym, K., Heitmann, K., Rabus, R., Schlesner, H., Amann, R., Reinhardt, R. (2003) Complete
 103 genome sequence of the marine planctomycete *Pirellula* sp. strain 1. Proc. Natl. Acad. Sci. USA 100,
 104 8298–8303.
 105
 106 Klindworth, A., Richter, M., Richter-Heitmann, T., Wegner, C.-E., Frank, C.S., Glöckner, F.O.
 107 Harder, J. (2014) Permanent draft genome of *Rhodopirellula rubra* SWK7. Mar. Gen. 13, 11-12.
 108
 109 Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W. (2014) Assessing the
 110 quality of microbial genomes recovered from isolates, single cells, and metagenomes. PeerJ PrePrints
 111 2:e554v1 <http://dx.doi.org/10.7287/peerj.preprints.554v1>.
 112
 113 Richter, M., Richter-Heitmann, T., Klindworth, A., Wegner, C.-E., Frank, C.S. Harder, J.,
 114 Glöckner, F.O. (2014a) Permanent draft genomes of the *Rhodopirellula maiorica* strain SM1. Mar.
 115 Gen. 13, 19-20.
 116
 117 Richter, M., Richter-Heitmann, T., Klindworth, A., Wegner, C.-E., Frank, C.S. Harder, J.,
 118 Glöckner, F.O. (2014b) Permanent draft genomes of the three *Rhodopirellula baltica* strains SH28,
 119 SWK14 and WH47. Mar. Gen. 13, 13-14.
 120
 121 Richter-Heitmann, T., Richter, M., Klindworth, A., Wegner, C.-E., Frank, C.S., Glöckner, F.O.
 122 Harder, J. (2014) Permanent draft genomes of the two *Rhodopirellula europaea* strains 6C and
 123 SH398. Mar. Gen. 13, 15-16.
 124
 125 Wegner, C.-E., Richter, M., Richter-Heitmann, T., Klindworth, A., Frank, C.S., Glöckner, F.O.

120 Harder, J. (2014) Permanent draft genome of *Rhodopirellula sallentina* SM41. Mar. Gen. 13, 17-18.

121 Winkelmann, N., Harder, J. (2009) An improved isolation method for attached-living *Planctomycetes*
122 of the genus *Rhodopirellula*. J. Microbiol. Meth. 77, 276-284.

123 Winkelmann, N., Jaekel, U., Meyer, C., Serrano, W., Rachel, R., Rosselló-Móra, R., Harder, J. (2010)
124 Determination of the diversity of *Rhodopirellula* isolates from European seas by multi-locus sequence
125 analysis. Appl. Envir. Microbiol. 76, 776-785.

126

127

128

129

130

131 Additional possible references:

132 Richter M., Rosselló-Móra, R. (2009) Shifting the genomic gold standard for the prokaryotic species
133 definition. Proc. Natl. Acad. Sci. USA 106, 19126-31.

134 Wegner, C.-E., Richter-Heitmann, T., Klindworth, A., Klockow, C., Richter, M., Achstetter, T., Glöckner, F.
135 O., Harder, J. (2013) Expression of sulfatases in *Rhodopirellula baltica* and the diversity of sulfatases in the
136 genus *Rhodopirellula*. Mar. Genom. 9, 51-61

137 Yilmaz, P et al., (2011) Minimum information about a marker gene sequence (MIMARKS) and minimum
138 information about any (x) sequence (MIXS) specifications. Nature Biotechnol. 29, 415-20.

139

140 Delcher, A.L., Bratke, K.A., Powers, E.C., Salzberg, E.C. (2007) Identifying bacterial genes and
141 endosymbiont DNA with Glimmer. *Bioinformatics* 23, 673–679.

142 Lagesen, K., Hallin, P., Rødland, E.A., Stærfeldt, H.-H., Rognes, T., Ussery, D.W. (2007) RNAmmer:
 143 Consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35, 3100-3108.

144 Lowe, T.M., and Eddy, S.R. (1997) tRNAscan-SE: a program for improved detection of transfer RNA genes
 145 in genomic sequence. *Nucleic Acids Res* 25, 955–964.

146 Meyer, F., Goesmann, A., McHardy, A.C., Bartels, D., Bekel, T., Clausen, J., et al. (2003) GenDB—an open
 147 source genome annotation system for prokaryote genomes. *Nucleic Acids Res.* 31, 2187–2195.

148 Mulder, N.J., Apweiler, R., Attwood, T.K., Bairoch, A., Bateman, A., Binns, D., et al. (2005) InterPro,
 149 progress and status in 2005. *Nucleic Acids Res.* 33, D201–D205.

150 Noguchi, H., Park, J., and Takagi, T. (2006) MetaGene: prokaryotic gene finding from environmental genome
 151 shotgun sequences. *Nucleic Acids Res.* 34, 5623–5630.

152 Quast, C. (2006) MicHanThi – design and implementation of a system for the prediction of gene functions in
 153 genome annotation projects. Master Thesis. Bremen, Germany: University of Bremen.

154 Richter, M., Lombardot, T., Kostadinov, I., Kottmann, R., Duhaime, M.B., Peplies, J., and Glöckner, F.O.
 155 (2008) JCoast – a biologist-centric software tool for data mining and comparison of prokaryotic metagenomes.
 156 *BMC Bioinforma.* 9, 177.

157

158 Schlesner, H., Rensmann, C., Tindall, B.J., Gade, D., Rabus, R., Pfeiffer, S., Hirsch, P. (2004) Taxonomic
 159 heterogeneity within the *Planctomycetales* as derived by DNA-DNA hybridization, description of
 160 *Rhodopirellula baltica* gen. nov., sp nov., transfer of *Pirellula marina* to the genus *Blastopirellula* gen. nov as
 161 *Blastopirellula marina* comb. nov and emended description of the genus *Pirellula*. *Int. J. Syst. Evol. Microbiol.*
 162 54, 1567-1580.

163